



THE MAN OF STEEL

SUPERHEROS, THEFT AND A WEIRD PROBLEM

STEPHEN E. HOHMANN, MD FACS

BAYLOR UNIVERSITY MEDICAL CENTER DALLAS



DISCLOSURES

- SPEAKER FOR MERIT MEDICAL
- SPEAKER FOR W.L. GORE

A SHOUT OUT




WHAT IS CLARK KENT'S MIDDLE NAME?

- A. ROBERT
- B. THOMAS
- C. WALTER
- D. JOSEPH
- E. WOLFGANG

WHAT SUPERVILLAIN SUPERMAN #75?

- A. MARK GLICKMAN
- B. JOHN ROSS
- C. DAVID CULL
- D. INGEMAR DAVIDSON
- E. DOOMSDAY



cull

verb

1.1.

select from a large quantity; obtain from a variety of sources.
"anecdotes culled from Greek and Roman history"

• *noun*

1.1.

a selective slaughter of wild animals.



WHEN DID LOIS AND SUPERMAN FIRST KISS?

- A. SUPERMAN #240
- B. ACTION COMICS #1
- C. SUPERMAN#3
- D. ACTION COMICS #78
- E. NEVER DO

WEDNESDAY AFTERNOON

- 50 year old female
- Left arm avg placed 3 months ago
- Gangrene at tip of index finger
- Finger pressure at rest 25mmHg and 60mmHg with compression
- Flow volume ~750cc/min



WHAT'S IN A NAME?

- STEEL – OR BETTER YET STEAL 😊
- DAIHS – DIALYSIS ACCESS INDUCED HAND ISCHEMIA
- DHIS – DISTAL HYPOPERFUSION ISCHEMIC SYNDROME
- DASS – DIALYSIS ACCESS STEAL SYNDROME

INCIDENCE

- 1-3% DISTAL RADIAL ARTERY AVFS
- 10-20% DISTAL BRACHIAL ARTERY AVFS
- HIGHER IN FEMORAL LOCATION
- SHOULD INCIDENCE BE 100% AND SYMPTOMATIC BE THE PERCENTAGE WE TALK ABOUT?



WHERE'S THE PROBLEM?

- ARTERIAL VASCULATURE WITH INTRINSIC DISEASE (PERIPHERAL ARTERIAL DISEASE)
- VERY HIGH FLOW IN AV ACCESS
- POOR COLLATERAL SUPPLY

PATHOPHYSIOLOGY

- PHYSIOLOGIC CHANGES WITH ACCESS CREATION
- BLOOD FLOW DIRECTION
 - PROXIMAL TO AV ACCESS: ALWAYS ANTEGRADE
 - DISTAL TO AV ACCESS: ANTEGRADE, BIPHASIC OR RETROGRADE
 - MAJORITY HAVE RETROGRADE FLOW
 - UPPER ARM AVGS 93%
 - RADIOCEPHALIICS 80%
 - BRACHIOCEPHALICS 73%
- ARTERIAL STENOSIS 1/3 DASS CASES



Stages of DASS

Stage 1

Palid/livid hand and/or cool hand without pain



Stage II

IIa - Tolerable pain during exercise and/or during dialysis

IIb - Intolerable pain during exercise and/or board during dialysis



Stage III

Pain at rest or loss of motor function



Stage IV

IVa - Limited tissue loss, potential for preservation of function



IVb - Irreversible tissue loss, significant loss of function

RISK FACTORS

TABLE V - PATIENTS AT RISK FOR STEAL

- Female
- Elderly (over 60 years of age)
- Diabetes
- Previous access surgeries in affected limb
- Use of distal brachial artery as anastomosis site
- Thumb pressures of 50 mmHg or less and pathological waveform, when occluding the ulnar artery

The Journal of Vascular Access 2008; 9: 155-166

Understanding the dialysis access steal syndrome. A review of the etiologies, diagnosis, prevention and treatment strategies

J. MALIK¹, V. TUKA¹, Z. KASALOVA¹, E. CHYTILOVA¹, M. SLAVIKOVA², P. CLAGETT³, I. DAVIDSON³, B. DOLMATCH⁴, D. NICHOLS⁵, M. GALLIEN⁶

SCREENING AND DIAGNOSIS

- HISTORY
- PHYSICAL EXAM
- ULTRASOUND INTERROGATION
- FINGER PRESSURES WITH AND WITHOUT COMPRESSION
- DBI (DIGITAL BRACHIAL INDEX) <0.6

TABLE I - STEAL SYMPTOMS IN ORDER OF INCREASING SEVERITY

- Nail changes
- Occasional tingling
- Extremity coolness
- Tingling and numbness in hand and fingers
- Muscle weakness
- Pale to whitish or cyanotic fingernail beds
- Rest-pain
- Sensory and motor function deficit
- Fingertip ulcerations
- Tissue loss

The Journal of Vascular Access 2008; 9: 155-166

Understanding the dialysis access steal syndrome. A review of the etiologies, diagnosis, prevention and treatment strategies

J. MALIK¹, V. TUKA¹, Z. KASALOVA¹, E. CHYTILOVA¹, M. SLAVIKOVA², P. CLAGETT³, I. DAVIDSON³, B. DOLMATECH⁴, D. NICHOLS⁵, M. GALLIEN⁶


TREATMENT OPTIONS

- OBSERVATION
- DISTAL RADIAL ARTERY LIGATION
- DISTAL REVASCULARIZATION INTERVAL LIGATION (DRIL)
- PROXIMAL ARTERIAL INFLOW (PAI)
- REVISION USING DISTAL INFLOW (RUDI)
- BANDING
- LIGATION

What about prevention?


DISTAL RADIAL ARTERY LIGATION

- LESS THAN 1% OF RADIOCEPHALIC AVFS
- OPEN OR ENDOVASCULAR
- BE SURE ARCH IS INTACT
- EXCELLENT RELIEF



DISTAL REVASCULARIZATION INTERVAL LIGATION

- SCHANZER 1988
- PROVIDING A “VERY LARGE COLLATERAL”
- GSV PREFERRED OVER PTFE
- DISTANCE FROM ANASTOMOSIS (10CM)
- TO LIGATE OR NOT TO LIGATE
- GOING OUT OF STYLE . . .
- SYMPTOM RELIEF ~90% WITH BYPASS PATENCY ~90%




The Journal of Vascular Access 2008; 9: 155-166

Understanding the dialysis access steal syndrome. A review of the etiologies, diagnosis, prevention and treatment strategies

J. MALIK¹, V. TUKA¹, Z. KASALOVA¹, E. CHYTILOVA¹, M. SLAVIKOVA², P. CLAGETT³, I. DAVIDSON⁴, B. DOLMAYCH⁵, D. NICHOLS⁵, M. GALLIEN⁶

PROXIMAL ARTERIAL INFLOW (PAI)

- CURRENT PROM KING/QUEEN OF OPEN REVISION
- NO LIGATION OF NATIVE SYSTEM
- USUALLY PROSTHETIC (?4-7MM GRAFT V 6 OR 7MM)
- INCREASING LENGTH OF CIRCUIT
- HOW PROXIMAL IS PROXIMAL




The Journal of Vascular Access 2008; 9: 155-166

Understanding the dialysis access steal syndrome. A review of the etiologies, diagnosis, prevention and treatment strategies

J. MALIK¹, V. TUKA¹, Z. KASALOVA¹, E. CHYTILOVA¹, M. SLAVIKOVA², P. CLAGETT³, I. DAVIDSON³, B. DOLMAYCH⁴, D. NICHOLS⁵, M. GALLIEN⁶


REVISION USING DISTAL INFLOW (RUDI)

- SUCCESSFUL IN 90% OF CASES, BUT AT A COST OF HIGHER THROMBOSIS? (22% IN ONE STUDY) - ?HYPERPLASIA AT NEW ANASTOMOSIS
- 2-3 CM DISTAL TO BIFURCATION
- NO LIGATION OF NATIVE ARTERY




LIGATION

- ISCHEMIC MONOMELIC NEUROPATHY
- RELIABLE RELIEF OF SYMPTOMS
- ?OFFERED TO ALL WITH STEAL
- BACK TO SQUARE ONE



BANDING

- REDUCE BLOOD FLOW (500-600CC/MIN AVF; 600-700CC/MIN AVG)
- INTRAOPERATIVE MONITORING (FINGER PRESSURES, ULTRASOUND VASCULAR ACCESS FLOW)
- AVF BAND NEAR ANASTOMOSIS
- AVG BAND IN THE MIDDLE



BANDING 2.0

The MILLER banding procedure is an effective method for treating dialysis-associated steal syndrome

Gregg A. Miller¹, Naveen Goel², Alexander Yevgeny Savransky⁴, Konstantin Khariton¹,

¹American Access Care of Brooklyn, Brooklyn, New York, USA; ²American Access Care of Queens, Queens, New York, USA; ³American Access Care of Queens, Queens, New York, USA; ⁴American Access Care of Queens, Queens, New York, USA

<http://www.kidney-international.org>

© 2010 International Society of Nephrology




Figure 5 | An inflated angioplasty balloon inside the access. Two parallel, made approximately 1-3 cm from the a




Figure 6 | A peri-access tunnel is dissected subcutaneously using Kelly clamp blunt dissection, and a 2-0 monofilament ligature of Prolene is pulled under the access.






Figure 8 | Following the Minimally Invasive Limited Ligation Endoluminal-Assisted Revision (MILLER) banding procedure, the resistance band redirects flow, improving distal perfusion, and alleviating symptoms.

WHAT TO DO?



- 50 year old female
- Left arm avg placed 3 months ago
- Gangrene at tip of index finger
- Finger pressure at rest 25mmHg and 60mmHg with compression
- Flow volume ~750cc/min



WHAT IF ARTERIAL ANASTOMOSIS IS ALREADY PROXIMAL?
HOW PROXIMAL IS PROXIMAL?



ISCHEMIC MONOMELIC NEUROPATHY

- DIFFUSE NEUROLOGIC DYSFUNCTION (ALL 3 NERVES) WITH EXCRUCIATING PAIN
- ELDERLY DIABETIC FEMALES
- WARM HAND WITH PALPABLE PULSE
- BRACHIAL ARTERY BASED AV ACCESS
- VASA NERVORUM BLOOD SUPPLY

TAKE HOME POINTS

- ALL VASCULAR DIALYSIS ACCESSES HAVE PHYSIOLOGIC STEAL THAT NEED NOT BE TREATED.
- CONSERVATIVE APPROACH IN STAGES I AND II A.
- LIGATION IS RARELY NEEDED AS THE OUTCOME OF OTHER OPTIONS IS QUITE GOOD AND SAFE.
- SYMPTOMATIC STEAL IS CHALLENGING BUT REQUIRES INTENSE ATTENTION UNTIL RESOLVED.
- STAGE II A MAY IMPROVE WITH SPECIFIC NON-SURGICAL MEASURES.
- STAGES II B, III AND IV REQUIRE INTERVENTION FIRST WITH ARTERIOGRAM PERFORMED TO EXCLUDE AND TREAT ARTERIAL PATHOLOGY PRESENT IN UP TO A THIRD OF DASS.
- WRIST AVF INDUCED DASS (<1%): LIGATE DISTAL RADIAL ARTERY WITH A PERMANENT SUTURE.
- BASED ON ACCESS BLOOD FLOW RATE, PAI IS FAVORED WITH < 800ML/MIN FOR AVFS

TAKE HOME POINTS

- **HIGHER FLOW** FAVORS A BANDING PROCEDURE BY OPEN SURGERY OR ENDOVASCULAR PROCEDURE..
- BANDING PROCEDURES SUCCESS DEPEND ON QUALITY OF **INTRAOPERATIVE GUIDANCE** WITH ULTRASOUND ACCESS VOLUME FLOW, FINGER PRESSURES OR PULSE OXIMETRY.
- THE **PAI PROCEDURE IS FAVORED OVER THE DRIL**, BECAUSE OF THE INCREASED INCIDENCE CARDIOVASCULAR MORBIDITY IN THE AGING DIALYSIS POPULATION MAKING THE DRIL DISTAL BYPASS ANASTOMOSIS HARD, RISKY AND OFTEN NOT POSSIBLE.
- **PREEMPTIVE** PAI OR RUDI PROCEDURES ELIMINATES STEAL IN THE HIGH RISK POPULATION AND ALSO PREVENTS ADDITIONAL PROCEDURES AS THESE PATIENTS HAVE LIMITED LONGEVITY.
- OPEN OR ENDOVASCULAR IS CHOSEN BASED ON **TEAM SKILL SETS, AND RESOURCES**.
- A COMPLETE ACCESS SURGEON IS TRAINED TO MOVE SEAMLESSLY BETWEEN OPEN AND ENDOVASCULAR DIALYSIS ACCESS PROCEDURES AS REQUIRED. THIS SKILL SET IS FAVORED IN MANAGEMENT OF DASS AS DIAGNOSTIC ARTERIOGRAM PRECEDES ALL INVASIVE TREATMENT OPTIONS AND IDEALLY PERFORMED IN THE SAME OR SETTING.



THANK YOU FOR YOUR ATTENTION!!

